

**GREAT HEARTS ACADEMY – ALLERGY ACTION PLAN** for the 2021/2022 SCHOOL YEAR

FIRST NAME: \_\_\_\_\_

LAST NAME: \_\_\_\_\_ DOB: \_\_\_\_\_

PARENT/GUARDIAN: \_\_\_\_\_

BEST CONTACT PHONE NUMBER: \_\_\_\_\_

PHYSICIAN NAME: \_\_\_\_\_

PHYSICIAN PHONE NUMBER: \_\_\_\_\_

TEACHER: \_\_\_\_\_ ROOM # \_\_\_\_\_

ALLERGIES: \_\_\_\_\_

TYPE OF REACTION: \_\_\_\_\_ Anaphylaxis \_\_\_\_\_ Nausea/Vomiting \_\_\_\_\_ Rash

Other reaction: \_\_\_\_\_

Allergic reaction may occur by: \_\_\_\_\_ Ingestion \_\_\_\_\_ Inhalation \_\_\_\_\_ Touch or Other: \_\_\_\_\_

Is the student asthmatic? \_\_\_\_\_ yes \_\_\_\_\_ no

My student will be eating food provided by local vendors for lunch \_\_\_\_\_ yes \_\_\_\_\_ no

My child may exhibit **MILD** symptoms with exposure to allergen \_\_\_\_\_

Treatment of **MILD** symptoms include:

1. Note time and occurrence of symptoms and stay with student
2. Watch closely for any sign of a serious reaction
3. Call parent/guardian listed above or communicate in writing of event
4. Give the following Medication: \_\_\_\_\_ Given to nurse \_\_\_\_\_ yes \_\_\_\_\_ date  
Dose: \_\_\_\_\_  
May repeat: \_\_\_\_\_  
Other instructions: \_\_\_\_\_
5. Call 911 or give emergency medications if symptoms worsen

My child may exhibit **SEVERE** symptoms with exposure to allergen \_\_\_\_\_

(Exhibiting any or all of the following symptoms is considered to be a severe allergic reaction: widespread hives and flushing, widespread tissue swelling, swelling of the tongue, throat itching or a sense of tightness in the throat, hoarseness and/or hacking cough, vomiting, nausea, cramps, diarrhea, repetitive coughing, wheezing, trouble breathing, rapid heart rate, lightheadedness, dizziness, loss of consciousness) Treatment of **SEVERE** symptoms include:

1. Note time and occurrence of symptoms and stay with student
2. Call 9-1-1 and inform them of a severe allergic reaction
3. Administer according to package instructions(circle) EpiPen 0.3 mg intramuscularly Given to nurse \_\_\_\_\_ yes  
EpiPen Jr. 0.15 mg intramuscularly  
Twinject 0.3 mg intramuscularly  
Twinject 0.15 mg intramuscularly
4. Call parent/guardian listed above, continue monitoring student for return of severe symptoms
5. Give injection device used, packaging, and student information to emergency responders
6. Give the following ANTIHISTAMINE: \_\_\_\_\_ Given to nurse \_\_\_\_\_ yes \_\_\_\_\_ date  
Dose: \_\_\_\_\_  
May repeat: \_\_\_\_\_  
Other instructions: \_\_\_\_\_

I understand that school staff **MUST** be informed of my child's health concerns in order to provide safe and appropriate care. I will update the school nurse office as my child's health conditions/treatments change throughout the year.

Parent/Guardian signature: \_\_\_\_\_ Date: \_\_\_\_\_

# Food Allergy Action Plan

## Emergency Care Plan

Place  
Student's  
Picture  
Here

Name: \_\_\_\_\_

D.O.B.: \_\_\_\_\_

Allergy To: \_\_\_\_\_

Weight: \_\_\_\_\_ lbs. Asthma:  Yes (higher risk for a severe reaction)  No

Any SEVERE SYMPTOMS after suspected or known ingestion:

One or more of the following:

LUNG: Short of breath, wheeze, repetitive cough

HEART: Pale, blue, faint, weak pulse, dizzy, confused

THROAT: Tight, hoarse, trouble breathing/swallowing

MOUTH: Obstructive swelling (tongue and/or lips)

SKIN: Many hives over body

Or combination of symptoms from different body areas:

SKIN: Hives, itchy rashes, swelling (e.g., eyes, lips)

GUT: Vomiting, diarrhea, crampy pain

1. **INJECT EPINEPHRINE IMMEDIATELY**

2. Call 911

3. Begin monitoring (see box below)

4. Give additional medications:\*

-Antihistamine

-Inhaler (bronchodilator) if asthma

\*Antihistamines & inhalers/bronchodilators are not to be depended upon to treat a severe reaction (anaphylaxis). USE EPINEPHRINE.

1. **GIVE ANTIHISTAMINE**

2. Stay with student; alert healthcare professionals and parent

3. If symptoms progress (see above), USE EPINEPHRINE

4. Begin monitoring (see box below)

### Medications/Doses

Epinephrine (brand and dose) \_\_\_\_\_

Antihistamine (brand and dose): \_\_\_\_\_

Other (e.g., inhaler-bronchodilator if asthmatic): \_\_\_\_\_

Parent/Guardian Signature \_\_\_\_\_


Date \_\_\_\_\_

Physician/Healthcare Provider Signature \_\_\_\_\_


Date \_\_\_\_\_

**EPIPEN Auto-Injector and  
EPIPEN Jr Auto-Injector Directions**

- First, remove the EPIPEN Auto-Injector from the plastic carrying case
- Pull off the blue safety release cap



- Hold orange tip near outer thigh (always apply to thigh)



- Swing and firmly push orange tip against outer thigh. Hold on thigh for approximately 10 seconds. Remove the EPIPEN Auto-Injector and massage the area for 10 more seconds



DEY™ and the Dey logo, EpiPen™, EpiPen 2-Pak®, and EpiPen Jr 2-Pak are registered trademarks of Dey Pharma, L.P.

**Adrenaclick™ 0.3 mg and  
Adrenaclick™ 0.15 mg Directions**



Remove GREY caps labeled "1" and "2."



Place RED rounded tip against outer thigh press down hard until needle penetrates. Hold for 10 seconds, then remove.

A food allergy response kit should contain at least two doses of epinephrine, other medications as noted by the student's physician, and a copy of this Food Allergy Action Plan.

A kit must accompany the student if he/she is off school grounds (i.e., field trip).

**Contacts:**

Call to 911: \_\_\_\_\_

Doctor: \_\_\_\_\_

Phone: \_\_\_\_\_

Parent/Guardian: \_\_\_\_\_

Phone: \_\_\_\_\_

Name/Relationship: \_\_\_\_\_

Phone: \_\_\_\_\_

